

A Nonstandard Numerical Scheme for a Predator-Prey Model Involving Predator Cannibalism and Refuge

Maya Rayungsari^{1,2}, Agus Suryanto^{1*}, W. M. Kusumawinahyu¹, Isnani Darti¹

¹Department of Mathematics, Faculty of Mathematics and Natural Sciences, University of Brawijaya, Malang 65145, Indonesia

²Department of Mathematics Education, Faculty of Pedagogy and Psychology, PGRI Wiranegara University, Pasuruan 67118, Indonesia

*Email: suryanto@ub.ac.id

Abstract

In this study, we implement a Nonstandard Finite Difference (NSFD) scheme for a predator-prey model involving cannibalism and refuge in predator. The scheme which is considered as a discrete dynamical system is analyzed. The performed analysis includes the determination of equilibrium point and its local stability. The system has four equilibrium points, namely the origin, the prey extinction point, the predator extinction point, and the coexistence point, which have exactly the same form and existence conditions as those in continuous system. The local stability of each first three equilibrium points is consistent with the one in continuous system. The stability of the coexistence point depends on the integration time step size. Nevertheless, the NSFD scheme allows us to choose the integration time step size for the solution to converge to a feasible point more flexible than the Euler and 4th order Runge-Kutta schemes. These are shown via numerical simulations.

Keywords: nonstandard finite difference scheme, predator-prey model, cannibalism, refuge

2010 MSC classification number: 70G60, 70Kxx, 92Bxx, 93C55, 92D25

1. INTRODUCTION

In recent decades, mathematical modelling of predator-prey interaction has attracted enormous interest from researchers and has been rapidly developed. Among the developments may applied the assumptions of the existing of harvesting [17], [24], [13], spread of disease [15], [11], Allee effect [16], [27], and stage-structure [5], [26]. Another interesting development is by adding the assumption of cannibalism existence [19]. The following model is a predator-prey model involving cannibalism in predator proposed by Deng et al [8].

$$\begin{aligned}\frac{dN}{dt} &= rN \left(1 - \frac{N}{K}\right) - b_1NP, \\ \frac{dP}{dt} &= c_1NP + c_2P - eP - \frac{b_2P^2}{k_2 + P}.\end{aligned}\tag{1}$$

$N \geq 0$ and $P \geq 0$ in (1) are the size population of prey and predator, respectively. In the absence of predator, prey grows logistically with a rate of r and the environment's carrying capacity of K . Then, in the presence of predator, prey decreases by b_1NP which is a type I Holling functional response with the predation rate of b_1 . Prey biomass consumed by predator is converted into predator births at a rate of c_1 . The predator dies naturally at a rate of e and decreases due to cannibalism by $\frac{b_2P^2}{k_2 + P}$, with b_2 and k_2 denote the maximum cannibalism rate and half-saturation constant of cannibalism, respectively. The cannibalism is converted to predator birth linearly at a rate of c_2 .

*Corresponding author

Received January 3rd, 2023, Revised February 25th, 2023 (first), Revised April 10th, 2023 (second), Accepted for publication May 5th, 2023. Copyright ©2023 Published by Indonesian Biomathematical Society, e-ISSN: 2549-2896, DOI:10.5614/cbms.2023.6.1.2

The system (1) had been modified by changing the predation rate following the type II Holling functional response and adding the assumption of the cannibalized predator refuge [18]. The modified system is

$$\begin{aligned}\frac{dN}{dt} &= rN \left(1 - \frac{N}{K}\right) - \frac{b_1 NP}{k_1 + N}, \\ \frac{dP}{dt} &= \frac{c_1 NP}{k_1 + N} + c_2 P - eP - \frac{b_2(1-m)P^2}{k_2 + (1-m)P},\end{aligned}\quad (2)$$

with k_1 denotes half-saturation constant of predation and m is proportion of refuge, while b_1 becomes the maximum predation rate. The dynamical properties of (2), those are the existence of equilibrium points, local and global stability analysis of the equilibrium points, and forward bifurcation existence had been shown in [18].

The predator-prey model (2) is a continuous dynamic system which is in the form of a system of nonlinear differential equations. Meanwhile, in general, nonlinear differential equations system cannot be solved analytically. Hence, discretization is required for approximating their solutions [3]. Moreover, as the stability of the equilibrium points in discrete models typically depends on the integration time stepsize, the discrete model mostly results in richer dynamics than the continuous one [9], [25].

A discrete model of a system of first-order differential equations can be obtained by applying the discretization method. Euler's method is one of the simplest method to discretize systems of differential equations. However, Mickens [12] had shown that the Euler discretization method has a weakness, that is the numerical instability. To eliminate it, Mickens proposes another method, namely the Nonstandard Finite Different (NSFD) scheme. By adopting the exact finite difference scheme, NSFD becomes a generalization of the Euler scheme. The most important property of a NSFD is that, in most cases, it eliminates the instability of the Euler scheme. NSFD follows the rules described below [12].

- 1) The first derivative of the differential equations system is approximated by the generalization of Euler's forward difference scheme, i.e

$$\frac{dX(n)}{dt} \rightarrow \frac{X(n+1) - X(n)}{\phi},$$

with $X(n) = X(t(n))$, $\phi = \phi(h)$ is a denominator function such that $\phi(h) = h + O(h^2)$, and h is the integration time step size.

- 2) The nonlinear form must be replaced with a non-local form like the following examples.

$$\begin{aligned}X^2 &\rightarrow X(n+1)X(n), \\ XY &\rightarrow \frac{X(n+1) + X(n-1)}{2}Y(n), \text{ etc.}\end{aligned}$$

If a discrete model and its corresponding continuous model have the same dynamic properties, they are said to be dynamically consistent [2]. The consistency of dynamics can be obtained using the discretization method of NFDS. NSFD has also been applied by researchers recently to develop biomathematics models into discrete systems [21], [6], [10], [1], [23], [7], [4], [14]. Therefore, in this paper, a NSFD scheme used to study discrete dynamical system of predator-prey model involving cannibalism and refuge in predator.

2. MODEL DEVELOPMENT

The continuous differential equation system (2) is discretized as follows.

$$\begin{aligned}N^2 &\rightarrow N(n+1)N(n), \\ NP &\rightarrow N(n+1)P(n), \\ P^2 &\rightarrow P(n+1)P(n), \\ \frac{dN}{dt} &\rightarrow \frac{N(n+1) - N(n)}{\phi(h)}, \\ \frac{dP}{dt} &\rightarrow \frac{P(n+1) - P(n)}{\phi(h)},\end{aligned}\quad (3)$$

where the denominator function $\phi(h)$ is $\phi(h) = h$. Through the discretization, the nonstandard numerical scheme for the system (2) is

$$\begin{aligned} N(n+1) &= N(n) + h \left(rN(n) \left(1 - \frac{N(n+1)}{K} \right) - \frac{b_1 N(n+1)P(n)}{k_1 + N(n)} \right), \\ P(n+1) &= P(n) + h \left(\frac{c_1 N(n+1)P(n)}{k_1 + N(n)} + c_2 P(n) - eP(n+1) - \frac{b_2(1-m)P(n+1)P(n)}{k_2 + (1-m)P(n)} \right). \end{aligned} \quad (4)$$

By using some algebraic manipulations, we get the explicit form for System (4), i.e.

$$\begin{aligned} N(n+1) &= \frac{N(n)(1+hr)}{\left[1 + h \left(\frac{rN(n)}{K} + \frac{b_1 P(n)}{k_1 + N(n)} \right) \right]} \equiv G_1(N(n), P(n)), \\ P(n+1) &= \frac{P(n)}{\left[1 + h \left(e + \frac{b_2(1-m)P(n)}{k_2 + (1-m)P(n)} \right) \right]} \left(1 + \frac{hc_1 N(n)(1+hr)}{(k_1 + N(n)) \left[1 + h \left(\frac{rN(n)}{K} + \frac{b_1 P(n)}{k_1 + N(n)} \right) \right]} + hc_2 \right) \equiv G_2(N(n), P(n)). \end{aligned} \quad (5)$$

To investigate the local stability of the equilibrium points of the system (3), the following lemma is very useful.

Lemma 2.1. [22] Consider $E^* = (N^*, P^*)$ is an equilibrium point of a two dimensional discrete system. For the quadratic equation $\lambda^2 - \tau(J(E^*))\lambda + \Delta(J(E^*)) = 0$, the roots satisfy $|\lambda_i| < 1, \forall i = 1, 2$, if and only if all of the following conditions are satisfied.

- 1) $1 + \tau(J(E^*)) + \Delta(J(E^*)) > 0$;
- 2) $1 - \tau(J(E^*)) + \Delta(J(E^*)) > 0$;
- 3) $\Delta(J(E^*)) < 1$;

where $\tau(J(E^*))$ and $\Delta(J(E^*))$, respectively, are the trace and determinant of characteristic equation of the Jacobian matrix at E^* .

3. LOCAL STABILITY

The equilibrium points of system (3) are obtained by finding the solutions of

$$\begin{aligned} G_1(N(n), P(n)) &= N(n), \\ G_2(N(n), P(n)) &= P(n), \end{aligned}$$

which is equivalent to the following system.

$$\begin{aligned} rN^* \left(1 - \frac{N^*}{K} \right) - \frac{b_1 N^* P^*}{k_1 + N^*} &= 0, \\ \frac{c_1 N^* P^*}{k_1 + N^*} + c_2 P^* - eP^* - \frac{b_2(1-m)P^{*2}}{k_2 + (1-m)P^*} &= 0. \end{aligned}$$

The equilibrium points are:

- 1) The origin, $E_0(0, 0)$, that clearly exists in $\mathbb{R}_+^2 = \{(N, P) : N \geq 0, P \geq 0\}$.
- 2) The prey extinction point, $E_1 \left(0, \frac{k_2(e - c_2)}{(c_2 - e - b_2)(1 - m)} \right)$, that exists in \mathbb{R}_+^2 if $0 < c_2 - e < b_2$.
- 3) The predator extinction point, $E_2(K, 0)$. E_2 always exists in \mathbb{R}_+^2 since $K > 0$.
- 4) The coexistence point, $E_3(N_3, P_3)$.

If $b_2 + e \neq c_1 + c_2$, N_3 and P_3 are derived by Cardanos's formula in [20] i.e

$$\begin{aligned} N_3 &= \frac{\sqrt[3]{Q_2 \pm \sqrt{Q_2^2 + \frac{4}{27}Q_1^3}}}{\sqrt[3]{2}} - \frac{Q_1 \sqrt[3]{2}}{3 \sqrt[3]{Q_2 \pm \sqrt{Q_2^2 + \frac{4}{27}Q_1^3}}} - \frac{B}{3A}, \\ P_3 &= \frac{r}{b_1} \left(1 - \frac{N_3}{K} \right) (k_1 + N_3), \end{aligned} \quad (6)$$

with

$$\begin{aligned}
Q_1 &= \frac{3AC - B^2}{3A^2}, \\
Q_2 &= \frac{9ABC - 2B^3 - 27A^2D}{27A^3}, \\
A &= \frac{r}{b_1K}(1-m)(b_2 - c_1 - c_2 + e), \\
B &= \frac{r}{b_1}(1-m) \left[(c_1 + c_2 - e - b_2) - \frac{k_1}{K}(c_1 + 2(c_2 - e - b_2)) \right], \\
C &= (c_1 + c_2 - e)k_2 + \frac{rk_1}{b_1}(1-m) \left[c_1 + (2 - k_1)(c_2 - e) - 2b_2 + \frac{b_2k_1}{K} \right], \text{ and} \\
D &= k_1 \left[k_2(c_2 - e) + \frac{rk_1}{b_1}(1-m)(c_2 - e - b_2) \right].
\end{aligned} \tag{7}$$

The point (6) exists in \mathbb{R}_+^2 if

- a) $Q_2^2 + \frac{4}{27}Q_1^3 \geq 0$ and
- b) $0 < N_3 < K$.

For $b_2 + e = c_1 + c_2$, the value of N_3 and P_3 are as follows.

$$N_3 = \frac{-C \pm \sqrt{C^2 - 4BD}}{2B}, P_3 = \frac{r}{b_1} \left(1 - \frac{N_3}{K} \right) (k_1 + N_3), \tag{8}$$

with

$$\begin{aligned}
B &= \frac{c_1rk_1}{b_1K}(1-m), \\
C &= b_2k_2 + \frac{rk_1}{b_1}(1-m) \left(k_1(c_1 - b_2) - c_1 + \frac{b_2k_1}{K} \right), \text{ and} \\
D &= k_1 \left[k_2(b_2 - c_1) - \frac{rc_1k_1}{b_1}(1-m) \right].
\end{aligned} \tag{9}$$

The coexistence point (8) exists in \mathbb{R}_+^2 if

- a) $C^2 - 4BD \geq 0$ and
- b) $0 < N_3 < K$.

It is found that the discrete system (3) has the exactly same equilibrium points as those in continuous model studied by Rayungsari et al [18].

For any equilibrium point of System (3), E^* , linearization of System (3) around E^* yield the Jacobian matrix

$$J(E^*) = \begin{bmatrix} \frac{\partial G_1}{\partial N} & \frac{\partial G_1}{\partial P} \\ \frac{\partial G_2}{\partial N} & \frac{\partial G_2}{\partial P} \end{bmatrix}_{E^*}, \tag{10}$$

where

$$\begin{aligned}
\frac{\partial G_1}{\partial N} &= \frac{1+hr}{\left[1+h\left(\frac{rN^*}{K} + \frac{b_1P^*}{k_1+N^*}\right)\right]^2} \left(1 + \frac{hb_1(k_1+2N^*)P^*}{(k_1+N^*)^2}\right), \\
\frac{\partial G_1}{\partial P} &= -\frac{(1+hr)N^*}{\left[1+h\left(\frac{rN^*}{K} + \frac{b_1P^*}{k_1+N^*}\right)\right]^2} \left(\frac{hb_1}{k_1+N^*}\right), \\
\frac{\partial G_2}{\partial N} &= \frac{(1+hr)hc_1P^*}{\left[1+h\left(e + \frac{b_2(1-m)P^*}{k_2+(1-m)P^*}\right)\right]^2} \left(\frac{k_1(1+hr) + hrN^*\left[1 - \frac{k_1+2N^*}{K}\right]}{(k_1+N^*)^2 \left[1+h\left(\frac{rN^*}{K} + \frac{b_1P^*}{k_1+N^*}\right)\right]^2}\right), \\
\frac{\partial G_2}{\partial P} &= \frac{1+he + \left(\frac{hb_2(1-m)^2P^{*2}}{(k_2+(1-m)P^*)^2}\right)}{\left[1+h\left(e + \frac{b_2(1-m)P^*}{k_2+(1-m)P^*}\right)\right]^2} \left(1 + \frac{hc_1N^*(1+hr)}{(k_1+N^*) \left[1+h\left(\frac{rN^*}{K} + \frac{b_1P^*}{k_1+N^*}\right)\right]} + hc_2\right) \\
&\quad - \frac{h^2b_1c_1N^*(1+hr)P^*}{\left[1+he + \frac{hb_2(1-m)P^*}{k_2+(1-m)P^*}\right] (k_1+N^*)^2 \left[1+h\left(\frac{rN^*}{K} + \frac{b_1P^*}{k_1+N^*}\right)\right]^2}.
\end{aligned} \tag{11}$$

The eigenvalues of the Jacobian matrix (10) are used to determine the local stability properties of the equilibrium points of system (3). The results are stated in Theorem 3.1.

Theorem 3.1. *The local stability properties of the equilibrium points of (3) are as follows.*

- 1) $E_0(0,0)$ is saddle if $c_2 < e$ and source if $c_2 > e$.
- 2) $E_1\left(0, \frac{k_2(e-c_2)}{(c_2-e-b_2)(1-m)}\right)$ is locally asymptotically stable if $r < \frac{b_1k_2(e-c_2)}{k_1(c_2-e-b_2)(1-m)}$ and saddle if $r > \frac{b_1k_2(e-c_2)}{k_1(c_2-e-b_2)(1-m)}$.
- 3) $E_2(K,0)$ is locally asymptotically stable if $c_1 < \frac{(e-c_2)(k_1+K)}{K}$ and saddle if $c_1 > \frac{(e-c_2)(k_1+K)}{K}$.
- 4) $E_3(N_3, P_3)$ is locally asymptotically stable if h satisfy all of the following conditions.
 - a) $1 + \tau(J(E_3)) + \Delta(J(E_3)) > 0$,
 - b) $1 - \tau(J(E_3)) + \Delta(J(E_3)) > 0$,
 - c) $\Delta(J(E_3)) - 1 < 0$,

with $\tau(J(E_3))$ and $\Delta(J(E_3))$ are the trace and determinant of characteristic equation of the Jacobian matrix (10) at E_3 , respectively. All of the three conditions will be computed numerically due to the terms' complexity.

Proof:

- 1) By substituting $E_0(0,0)$ to (10), we get

$$J(E_0) = \begin{bmatrix} 1+hr & 0 \\ 0 & \frac{1+hc_2}{1+he} \end{bmatrix}.$$

The eigenvalues of $J(E_0)$ are $\lambda_1 = 1+hr$ and $\lambda_2 = \frac{1+hc_2}{1+he}$. Since $|\lambda_1| > 1$, E_0 is unstable, more precisely saddle if $c_2 < e$ and source if $c_2 > e$.

- 2) The Jacobian matrix for prey extinction point is

$$J(E_1) = \begin{bmatrix} \frac{1+hr}{1+\frac{hb_1P_1}{k_1}} & 0 \\ \frac{(1+hr)hc_1P_1}{\left[1+h\left(e + \frac{b_2(1-m)P_1}{k_2+(1-m)P_1}\right)\right]^2} k_1 \left(1 + \frac{hb_1P_1}{k_1}\right)^2 & \frac{1+he + \left(\frac{hb_2(1-m)^2P_1^2}{(k_2+(1-m)P_1)^2}\right)}{\left[1+h\left(e + \frac{b_2(1-m)P_1}{k_2+(1-m)P_1}\right)\right]^2} (1+hc_2) \end{bmatrix},$$

with $P_1 = \frac{k_2(e - c_2)}{(c_2 - e - b_2)(1 - m)}$. $J(E_1)$ has eigenvalues $\lambda_1 = \frac{1 + hr}{1 + \frac{hb_1 P_1}{k_1}} = \frac{1 + hr}{1 + \frac{hb_1 k_2(e - c_2)}{k_1(c_2 - e - b_2)(1 - m)}}$
 and $\lambda_2 = \frac{1 + he + \left(\frac{hb_2(1 - m)^2 P_1^2}{(k_2 + (1 - m)P_1)^2}\right)}{\left[1 + h\left(e + \frac{b_2(1 - m)P_1}{k_2 + (1 - m)P_1}\right)\right]^2} (1 + hc_2)$. Through some algebraic manipulations, we find that

$$\lambda_2 = \frac{1 + he + \left(\frac{hb_2(1 - m)^2 P_1^2}{(k_2 + (1 - m)P_1)^2}\right)}{\left[1 + h\left(e + \frac{b_2(1 - m)P_1}{k_2 + (1 - m)P_1}\right)\right]^2} (1 + hc_2) = \frac{1 + h\left(\frac{e + (e - c_2)^2}{b_2}\right)}{1 + hc_2}.$$

The existence conditions of E_1 , $0 < c_2 - e < b_2$, leads to $e + \frac{(e - c_2)^2}{b_2} < c_2$ and thus $|\lambda_2| < 1$.

If $r < \frac{b_1 k_2(e - c_2)}{k_1(c_2 - e - b_2)(1 - m)}$ then $|\lambda_1| < 1$ and E_1 is locally asymptotically stable. While, if $r > \frac{b_1 k_2(e - c_2)}{k_1(c_2 - e - b_2)(1 - m)}$ then $|\lambda_1| > 1$ and E_1 is saddle.

3) The Jacobian matrix for E_2 is

$$J(E_2) = \begin{bmatrix} \frac{1}{1 + hr} & -\frac{hb_1 K}{(1 + hr)(k_1 + K)} \\ 0 & \frac{1 + h\left(\frac{c_1 K}{k_1 + K} + c_2\right)}{1 + he} \end{bmatrix},$$

so that the eigenvalues are $\lambda_1 = \frac{1}{1 + hr}$ and $\lambda_2 = \frac{1 + h\left(\frac{c_1 K}{k_1 + K} + c_2\right)}{1 + he}$. It is clear that $|\lambda_1| < 1$.

If $c_1 < \frac{(e - c_2)(k_1 + K)}{K}$, then $|\lambda_2| < 1$ and E_2 locally asymptotically stable. Otherwise, if $c_1 > \frac{(e - c_2)(k_1 + K)}{K}$ and E_2 becomes saddle.

4) The Jacobian matrix for coexistence point is

$$J(E_3) = \begin{bmatrix} J_{11} & J_{12} \\ J_{21} & J_{22} \end{bmatrix},$$

where

$$\begin{aligned} J_{11} &= \frac{1 + hr}{[1 + hr]^2} \left(1 + \frac{hb_1(k_1 + 2N_3)P_3}{(k_1 + N_3)^2}\right) = \frac{1}{1 + hr} \left(1 + \frac{hb_1(k_1 + 2N_3)P_3}{(k_1 + N_3)^2}\right), \\ J_{12} &= -\frac{(1 + hr)N_3}{[1 + hr]^2} \left(\frac{hb_1}{k_1 + N_3}\right) = -\frac{N_3}{1 + hr} \left(\frac{hb_1}{k_1 + N_3}\right), \\ J_{21} &= \frac{hc_1 P_3}{\left[1 + \frac{hc_1 N_3}{k_1 + N_3} + hc_2\right]} \left(\frac{(k_1 + N_3)[1 + hr] - N_3 \left[1 + \frac{hr}{K}(k_1 + 2N_3)\right]}{(k_1 + N_3)^2(1 + hr)}\right), \\ J_{22} &= \frac{1 + he + \left(\frac{hb_2(1 - m)^2 P_3^2}{(k_2 + (1 - m)P_3)^2}\right)}{\left[1 + \frac{hc_1 N_3}{k_1 + N_3} + hc_2\right]} - \frac{h^2 r c_1 N_3 \left(1 - \frac{N_3}{K}\right)}{\left[1 + \frac{hc_1 N_3}{k_1 + N_3} + hc_2\right] (k_1 + N_3)(1 + hr)}. \end{aligned} \quad (12)$$

The eigenvalues of $J(E_3)$ are the roots of the characteristic function

$$\lambda^2 - \text{tr}(J(E_3))\lambda + \Delta(J(E_3)) = 0, \quad (13)$$

with $\text{tr}(J(E_3)) = J_{11} + J_{22}$ and $\Delta(J(E_3)) = J_{11}J_{22} - J_{12}J_{21}$. According to Lemma 2.1, E_3 is locally asymptotically stable if all of the following three conditions are satisfied.

$$\begin{aligned} (i) \quad & 1 + \tau(J(E_3)) + \Delta(J(E_3)) > 0, \\ (ii) \quad & 1 - \tau(J(E_3)) + \Delta(J(E_3)) > 0, \\ (iii) \quad & \Delta(J(E_3)) - 1 < 0. \end{aligned} \quad (14)$$

■

4. NUMERICAL SIMULATIONS

Numerical simulations of the system (3) are performed in this section via Matlab software. We apply three scheme, i.e NSFD, Euler, and 4th order Runge-Kutta. The numerical simulations aim to compare the consistency of system stability using the three methods. The parameter values used in the numerical simulations in this paper are assumed hypothetically since there is no available real data.

Table 1: Parameter Values.

Parameter	Simulation 1	Simulation 2	Simulation 3	Simulation 4
r	1	1	1	1
K	1	1	1	1
b_1	0.3	0.5	0.5	0.5
k_1	0.3	0.3	0.3	0.3
c_1	0.2	0.1	0.3	0.45
c_2	0.12	0.2	0.2	0.2
e	0.02	0.3	0.3	0.3
b_2	0.2	0.3	0.3	0.3
m	0.3	0.3	0.3	0.3
k_2	1	1	1	1

4.1. Simulation 1

By using the parameter values as in Table 1 (Simulation 1), the prey extinction point exists, i.e. $E_1(0, 1.4286)$. The stability condition of E_1 in Theorem 3.1 point (ii) is satisfied. It can be seen in Figure 1 that with a very small step size, $h = 0.05$, the system solutions using three methods convergent to E_1 . This is consistent with the results obtained in [18]. By increasing the step size to $h = 12$, the solutions of system (3) by NFSD remains convergent to E_1 . Meanwhile, the solutions obtained by Euler and 4th order Runge-Kutta methods can be negative (see Figure 2), which are not reliable.




Figure 1: Phase portraits of simulation 1 with $h = 0.05$ using: (a) NSFD, (b) Euler, and (c) 4th order Runge Kutta methods.

4.2. Simulation 2

To show the behavior of the solutions regarding the stability of predator extinction point, E_2 , the parameter values in Table 1 (Simulation 2) are chosen. With those parameter values, the stability conditons of $E_2(1, 0)$ are hold. This is in accordance with the results of the numerical simulation shown in Figure 3, with the step size used is $h = 0.05$. When the step size used is quite large as shown in Figure 4, i.e. $h = 3.6$, and by taking initial values near E_2 , the solutions of system (3) by Euler method is nonfeasible since they convergent to negative solutions, while the solutions by Runge Kutta leads to a periodic solutions of period two. Whereas, the stability of E_2 is still consistent for NFSD scheme.




Figure 2: Phase portraits of simulation 1 with $h = 12$ using: (a) NSFD, (b) Euler, and (c) 4th order Runge Kutta methods.




Figure 3: Phase portraits in simulation 2 with $h = 0.05$ using: (a) NSFD, (b) Euler, and (c) 4th order Runge Kutta methods.




Figure 4: Phase portraits in simulation 2 with $h = 3.6$ using: (a) NSFD, (b) Euler, and (c) 4th order Runge Kutta methods.

4.3. Simulation 3

With the parameter values in Table 1 (Simulation 3), the coexistence point exists, i.e. $E_3(0.6026, 0.7174)$. For any values of h , all of the stability conditions of E_3 in (14) are satisfied, as we can see in Figure 5. With a step size of $h = 0.05$ (see Figure 6), E_3 is asymptotically stable for the NSFD, Euler, and 4th order

Runge Kutta methods. By increasing the step size value to $h = 7.5$ (see Figure 7), with initial values close to E_3 , the solutions of system (2) using Euler and 4th order Runge Kutta methods lead to nonfeasible solutions (negative solutions), while the system with NSFD scheme is still consistent (see Figure 6). In Figure 7(c), the solution for predator goes down first then up.




Figure 5: Graphics of (a) $1 + \tau(J(E_3)) + \Delta(J(E_3))$, (b) $1 - \tau(J(E_3)) + \Delta(J(E_3))$, and (c) $\Delta(J(E_3)) - 1$, in simulation 3.




Figure 6: Phase portraits in simulation 3 with $h = 0.05$ using: (a) NSFD, (b) Euler, and (c) 4th order Runge Kutta methods.

4.4. Simulation 4

By increasing the value of c_1 , as in Table 1 (Simulation 4), the coexistence point is $E_3(0.2631, 0.8299)$. The first two stability conditions of E_3 in (14) are hold for any h , but the last condition isn't satisfied for $0 < h < 0.1006$ (see Figure 8(c)) and is satisfied for all $h \geq 0.1006$ (see Figure 8(d)). This is supported by the bifurcation diagram in Figure 9. For $h < 0.1006$, demonstrated particularly by $h = 0.05$ (Figure 10), the system convergent to a periodic solution. Meanwhile, for $h > 0.1006$ which is represented by $h = 12$ in Figure 11, the system convergent to coexistence point. In the continuous model in [18], with parameter values as in Table 1 (Simulation 4), the solutions do not converge to the coexistence point but to a limit cycle around it. In contrary, in the discrete model in this article, the solutions of system (3) can be stable to the coexistence point, by choosing the integration step-size larger than 0.1006.

Next, to compare the behavior of the system with the NSFD, Euler, and 4th order Runge Kutta schemes, we choose $h = 0.05$ in simulation illustrated by Figure 12. While, in Figure 13 we choose $h = 12$. We can see that in Figure 12, the solutions of the three schemes convergent to a limit cycle around E_3 . In Figure 13 ($h = 12$), the solutions of NSFD scheme convergent to E_3 , whereas the two other schemes with initial value close to E_3 convergent to nonfeasible (negative) solutions. In the results obtained in [18], the stability




Figure 7: Phase portraits in simulation 3 with $h = 7.5$ using: (a) NSFD, (b) Euler, and (c) 4th order Runge Kutta methods.




Figure 8: Graphics of (a) $1 + \tau(J(E_3)) + \Delta(J(E_3))$, (b) $1 - \tau(J(E_3)) + \Delta(J(E_3))$, (c) $\Delta(J(E_3)) - 1$ (for relative small h), and (d) $\Delta(J(E_3)) - 1$ (for $0 < h \leq 20$), in simulation 4.

of E_3 is the same as the results in Figure 12. Thus, in this case, the stability properties of the NSFD system are only consistent with the continuous system for very small values of h . However, the advantage of NSFD here is that it keeps the solution positive, unlike the Euler and 4th-order Runge-Kutta schemes which make the solutions turn to negative for large h values.




Figure 9: Diagram bifurcation in simulation 4 with h as bifurcation parameter: (a) $h - N^*$, (b) $h - P^*$.




Figure 10: Solution of N and P in simulation 4 with $h = 0.05$: (a) $t - N$, (b) $t - P$.




Figure 11: Solution of N and P in simulation 4 with $h = 12$: (a) $t - N$, (b) $t - P$.

5. CONCLUSION

In this paper, a discrete system of predator-prey model involving cannibalism and refuge in predator had been developed using Nonstandard Finite Difference (NFSD) scheme. Based on the analysis results, the system has four equilibrium points, i.e the origin, the prey extinction point, the predator extinction point, and the coexistence point which have exactly the same form and existence conditions as the ones in the continuous system. Furthermore, the local stability conditions for the first three equilibrium points have been




Figure 12: Phase portraits in simulation 4 with $h = 0.05$ using: (a) NSFD, (b) Euler, and (c) 4^{th} order Runge Kutta methods.




Figure 13: Phase portraits in simulation 4 with $h = 12$ using: (a) NSFD, (b) Euler, and (c) 4^{th} order Runge Kutta methods.

shown to be exactly the same as those in the continuous system. Meanwhile, there are differences in the coexistence point's stability conditions. In the discrete system using an NSFD scheme the stability depends on the integration time step size, which does not occur in the continuous system. This is confirmed in numerical simulations which show that in some cases, the stability of the coexistence point is only consistent with the one in continuous system at a relative small step size, and inconsistent for step size that exceeds the threshold. Nonetheless, unlike the Euler and 4^{th} order Runge-Kutta schemes, the NSFD scheme allows us to be more flexible in choosing the integration time stepsize for the solution to converge to a feasible point.

ACKNOWLEDGEMENT

This research is supported by Faculty of Mathematics and Natural Sciences (FMIPA) via Public Funds DPA (Dokumen Pelaksanaan Anggaran) Perguruan Tinggi Berbadan Hukum (PTNBH) University of Brawijaya and is based on FMIPA Professor Grant Contract No. 3084.19/UN10.F09/PN/2022.

REFERENCES

- [1] Adekanye, O. and Washington, T., Nonstandard finite difference scheme for a Tacoma Narrows Bridge model, *Applied Mathematical Modelling*, 62, pp. 223-236, 2018.
- [2] Al-Kahby, H., Dannan, F. and Elaydi, S., Non-standard discretization methods for some biological models, *Applications of nonstandard finite difference schemes*, pp. 155-180, 2000.
- [3] Anguelov, R. and Lubuma, J. M. S., Nonstandard finite difference method by nonlocal approximation, *Mathematics and Computers in Simulation*, 61(3-6), pp. 465-475, 2003.

- [4] Baleanu, D., Zibaei, S., Namjoo, M. and Jajarmi, A., A nonstandard finite difference scheme for the modeling and nonidentical synchronization of a novel fractional chaotic system, *Advances in Difference Equations*, 308(1), pp. 1-19, 2021.
- [5] Beay, L. K., Suryanto, A. and Darti, I., Hopf bifurcation and stability analysis of the Rosenzweig-MacArthur predator-prey model with stage-structure in prey, *Mathematical Biosciences and Engineering*, 17(4), pp. 4080-4097, 2020.
- [6] Darti, I. and Suryanto, A., Stability preserving non-standard finite difference scheme for a harvesting Leslie-Gower predator-prey model, *Journal of Difference Equations and Applications*, 21(6), pp. 528-534, 2015.
- [7] Darti, I. and Suryanto, A., Dynamics of a SIR epidemic model of childhood diseases with a saturated incidence rate: Continuous model and its nonstandard finite difference discretization, *Mathematics*, 8(9), p. 1459, 2020.
- [8] Deng, H., Chen, F., Zhu, Z. and Li, Z., Dynamic behaviors of Lotka-Volterra predator-prey model incorporating predator cannibalism, *Advances in Difference Equations*, 2019(1), pp. 1-17, 2019.
- [9] Elaydi, S., *An Introduction to Difference Equations*. Springer, San Antonio, Texas, 3rd edition, 2005.
- [10] Fitriah, Z., and Suryanto, A., Nonstandard finite difference scheme for SIRS epidemic model with disease-related death, *AIP Conference Proceedings*, 1723(1), p. 030009, 2016.
- [11] Gupta, A. and Dubey, B., Bifurcation and chaos in a delayed eco-epidemic model induced by prey configuration, *Chaos, Solitons, and Fractals*, 165, p. 112785, 2022.
- [12] Mickens, R. E., *Applications of Nonstandard Finite Difference Schemes*, World Scientific Publishing Co. Pte. Ltd., USA, 2000.
- [13] Panigoro, H. S., Rahmi, E., Achmad, N. and Mahmud, S. L., The influence of additive Allee effect and periodic harvesting to the dynamics of Leslie-Gower predator-prey model, *Jambura Journal of Mathematics*, 2(2), pp. 87-96, 2020.
- [14] Panigoro, H. S., Rayungsari, M. and Suryanto, A., Bifurcation and chaos in a discrete-time fractional-order logistic model with Allee effect and proportional harvesting, *International Journal of Dynamics and Control*, pp. 1-15, 2023.
- [15] Panigoro, H. S., Suryanto, A., Kusumawinahyu, W. M. and Darti, I., Dynamics of an eco-epidemic predator-prey model involving fractional derivatives with power-law and Mittag-Leffler kernel. *Symmetry*, 13(5), p. 785, 2021.
- [16] Rahmi, E., Darti, I., Suryanto, A. and Trisilowati, A modified Leslie-Gower model incorporating Beddington-DeAngelis functional response, double Allee effect and memory effect, *Fractal and Fractional*, 5(3), p. 84, 2021.
- [17] Rayungsari, M., Kusumawinahyu, W.M. and Marsudi, Dynamical Analysis of Predator-Prey Model with Ratio-Dependent Functional Response, *Applied Mathematical Sciences*, 8(29), pp. 1401-1410, 2014.
- [18] Rayungsari, M., Suryanto, A., Kusumawinahyu, W.M. and Darti, I., Dynamical analysis of a predator-prey model incorporating predator cannibalism and refuge, *Axioms*, 11(3), p. 116, 2022.
- [19] Rayungsari, M., Suryanto, A., Muharini, W. and Darti, I., Dynamics analysis of a predator-prey fractional-order model incorporating predator cannibalism and refuge, *Frontiers in Applied Mathematics and Statistics*, 9(29), pp. 1-12, 2023.
- [20] Ganti, S. and Gopinathan, S., A note on the solutions of cubic equations of state in low temperature region, *Journal of Molecular Liquids*, 315, p. 113808, 2020.
- [21] Suryanto, A., A nonstandard finite difference scheme for SIS epidemic model with delay: stability and bifurcation analysis, 2012.
- [22] Suryanto, A., Stability analysis of the Euler discretization for SIR epidemic model, In *AIP Conference Proceedings* 1602(1), pp. 375-379, American Institute of Physics, 2014.
- [23] Suryanto, A. and Darti, I., On the nonstandard numerical discretization of SIR epidemic model with a saturated incidence rate and vaccination, *AIMS Math*, 6, pp. 141-155, 2021.
- [24] Suryanto, A., Darti, I., S. Panigoro, H. and Kilicman, A., A fractional-order predator-prey model with ratio-dependent functional response and linear harvesting, *Mathematics*, 7(11), p. 1100, 2019.
- [25] Yue, D., Guan, Z.H., Chen, J., Ling, G. and Wu, Y., Bifurcations and chaos of a discrete-time model in genetic regulatory networks, *Nonlinear Dynamics*, 87(1), pp. 567-586, 2017.
- [26] Zhang, S., Yuan, S. and Zhang, T., A predator-prey model with different response functions to juvenile and adult prey in deterministic and stochastic environments, *Applied Mathematics and Computation*, 413, p. 126598, 2022.
- [27] Zhu, Z., Chen, Y., Li, Z. and Chen, F., Stability and bifurcation in a Leslie-Gower predator-prey model with Allee effect, *International Journal of Bifurcation and Chaos*, 32(03), p. 2250040, 2022.